

AD-A094 728

MASSACHUSETTS INST OF TECH LEXINGTON LINCOLN LAB F/G 12/1  
ON NONPARAMETRIC PROBABILITY DENSITY ESTIMATION USING ORTHOGONA--ETC(U)  
NOV 80 L K JONES F19628-80-C-0002

UNCLASSIFIED

TN-1980-54

ESD-TR-80-230

NL

[ref]  
AD 94 7-19



			END
			DATE
			FORMED
			3-8-81
			DTIC

AD 1009-1728

See 1473

LEVEL II

12

Technical Note

1980-54

On Nonparametric  
Probability Density Estimation  
Using Orthogonal Series

L. K. Jones


DTIC  
ELECTE  
FEB 9 1981  
S D E

26 November 1980

Prepared for the Department of the Air Force  
under Electronic Systems Division Contract F19628-80-C-0002 by

Lincoln Laboratory

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
LEXINGTON, MASSACHUSETTS



Approved for public release; distribution unlimited.

FILE COPY

81 2 09 00

2-29

The work reported in this document was performed at Lincoln Laboratory, a center for research operated by Massachusetts Institute of Technology, with the support of the Department of the Air Force under Contract F19628-60-C-0002.

This report may be reproduced to satisfy needs of U.S. Government agencies.

The views and conclusions contained in this document are those of the contractor and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the United States Government.

This technical report has been reviewed and is approved for publication.

FOR THE COMMANDER

*Raymond L. Lottelle*

Raymond L. Lottelle, Lt. Col., USAF  
Chief, ESD Lincoln Laboratory Project Office

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
LINCOLN LABORATORY

ON NONPARAMETRIC PROBABILITY DENSITY ESTIMATION  
USING ORTHOGONAL SERIES

L. K. JONES

Group 92

Accession For		
NTIS GRA&I	<input checked="checked" type="checkbox"/>	
DTIC TAB	<input type="checkbox"/>	
Unannounced	<input type="checkbox"/>	
Justification		
By		
Distribution		
Availability Codes		
Avail and/or		
Disc		
A		

TECHNICAL NOTE 1980-54

26 NOVEMBER 1980

Approved for public release; distribution unlimited.

LEXINGTON

MASSACHUSETTS

# ABSTRACT

A method of density estimation is proposed, which is a rational modification of orthogonal expansions, combined with a stopping rule determined by a nearest neighbor statistic. This method yields consistent estimates and applies (in principle) to density estimation in any number of dimensions.

# ON NONPARAMETRIC PROBABILITY DENSITY ESTIMATION USING ORTHOGONAL SERIES

## I. INTRODUCTION

Among the numerous non-parametric methods of estimating a probability density function, the approximation of this density by a finite fourier series has several computational advantages. Probably the most important of these is the fact that the evaluation of this density at a new data point requires only the storage of certain fourier coefficients. One of the main disadvantages of such an approximation of a density is the difficulty of determining the number of terms in the expansion.

In this note, we propose an approximation which is a rational function of a finite fourier series. The number of terms in this series depends, in a very natural way, on the nearest neighbor error rate for the sample data when compared to a sample drawn from a reference distribution. In III we show that the method is consistent and in IV we remark on the relevance of this method in hypothesis testing.

## II. SECOND ORDER SOLUTION TO THE BINARY DECISION PROBLEM

Let  $p_1, p_2$  be two Lebesgue measurable, bounded ( $\leq K$ ) probability density functions on the unit cube,  $I$ , in  $R^n$ . We assume further that  $p_1 \neq p_2$  on some set of positive measure in  $I$  and that for some  $\delta > 0$ ,  $p_1 \geq \delta$  on  $I$ . Let  $\mathcal{L} = \{f \in L_2(I) : E_1 f = \int f p_1 dx = 0, E_2 f = \int f p_2 dx = 1\}$ . According to [1], a second order solution for an optimal discriminant  $\bar{f} \in \mathcal{L}$ , for the binary hypothesis test  $H_1: X$  has density  $p_1$  vs.  $H_2: X$  has density  $p_2$ , is a critical point for some real  $\alpha$  of the functional

$$J_\alpha(f) = \alpha \text{VAR}_1 f + (1-\alpha) \text{VAR}_2 f \quad (1)$$

In fact if we restrict ourselves to the case  $0 < \alpha < 1$  and solve (1) for the unique (to within a null function) critical point (and global minimum of  $J_\alpha(f)$  for  $f \in \mathcal{L}$ ), we obtain by elementary variational calculus

$$\bar{f} = \frac{[(1-\alpha)-\lambda] p_2/p_1 + \lambda}{\alpha + (1-\alpha) p_2/p_1} \quad (2)$$

with

$$0 > \lambda = \frac{(1-\alpha) \int \frac{p_2 p_1}{\alpha p_1 + (1-\alpha) p_2} dx}{\int \frac{(p_2 - p_1) p_1}{\alpha p_1 + (1-\alpha) p_2} dx} = (1-\alpha) - J_\alpha(\bar{f})$$

It follows that  $\bar{f}$  is rational and increasing in  $(p_2/p_1)$  and hence optimal (by an adjustment of threshold) for minimum total error (or Neyman-Pearson at level  $\beta$ ) hypothesis testing.

### III. DENSITY ESTIMATION

For simplicity we consider only the case  $\alpha = \frac{1}{2}$ . Similar results may be obtained for other  $\alpha$ . Solving (2) for  $p_2/p_1$  we obtain

$$\frac{p_2}{p_1} = \frac{\frac{1}{2}\bar{f} - \lambda}{(\frac{1}{2} - \lambda) - \frac{1}{2}\bar{f}} = \frac{\bar{f} - 1 + 2J_{\frac{1}{2}}(\bar{f})}{2J_{\frac{1}{2}}(\bar{f}) - \bar{f}} \quad (3)$$

We now write

$$J_{\frac{1}{2}}(\bar{f}) = \frac{1}{2} + \frac{\epsilon_{nn}}{4(\frac{1}{2} - \epsilon_{nn})} \quad (4)$$

where  $\epsilon_{nn} = \int \frac{p_2 p_1}{p_1 + p_2} dx$  is known as the limiting nearest neighbor error rate, i.e., if we generate  $n$  independent class 1 samples from a distribution with density  $p_1$  and similarly  $n$  class 2 samples from a distribution with density  $p_2$  and then classify new samples (drawn from class 1 or class 2 with equal probability) as the class of the (a) nearest neighbor in the original  $2n$ , then the classification error of this procedure approaches  $\epsilon_{nn}$  as  $n \rightarrow \infty$  with probability 1. (See [2].)

We now make a final assumption that  $p_1 \equiv 1$  on  $I$ . Again, results analagous to the following will still hold provided  $p_1$  is strictly bounded away from 0 in  $I$ .

Suppose we are given  $n$  independent samples from a distribution with density  $p_2$ . Let  $1 = \varphi_1, \varphi_2, \dots$  be a complete orthonormal system for  $L_2(I)$ . Finally, let  $v_n$  be the empirical density determined by the  $n$  sample points. Now, consider the solution



of the variational problem: minimize

$$\frac{1}{2} \text{VAR}_1 f + \frac{1}{2} \text{VAR}_{V_n} f = J_N^n(f) \quad (*)$$

such that

$$f = \sum_{i=1}^N a_i \varphi_i$$

$$E_1 f = 0$$

$$E_{V_n} f = 1$$

where  $N$  is determined by a "stopping rule". We then let  $f_n$  be the above minimizing  $f$ . Before describing the determination of  $N$ , we show that the preceding variational problem has solutions with probability 1 for large enough  $N$ .

Lemma Assume  $n$  is fixed. Then with probability one  $(*)$  has solutions for large enough  $N$ . In fact  $\min_f J_N^n(f) \rightarrow 0$  as  $N \rightarrow \infty$  with probability one.

Proof: Let  $L$  be any positive integer and  $\epsilon > 0$ . Then there is an  $N_0$  such that, for  $N > N_0$ , there are functions  $\psi_1, \psi_2, \dots, \psi_L \in \langle \varphi_1, \varphi_2, \dots, \varphi_N \rangle$  with the properties:

$$(i) \quad \|\psi_i\|_2^2 \leq \frac{1}{L} + \epsilon$$

(ii) there exist disjoint subsets  $A_1, \dots, A_L$  with

$$m\left(\bigcup_{i=1}^L A_i\right) > 1 - \epsilon \quad \text{s.t. } x \in A_i \text{ implies}$$

$$|\psi_i(x) - 1| < \epsilon \quad \text{and} \quad |\psi_j(x)| < \epsilon \quad (j \neq i).$$

Hence, with probability (wrt  $p_2$ )  $> (1-K\epsilon)^n$ , each of our samples  $\ell$  will lie in some  $A_{i_\ell}$ . Let us now consider the function

$$\tilde{f} = \frac{\sum_{\ell=1}^n \psi_{i_\ell} - \sum_{\ell=1}^n \int \psi_{i_\ell} dx}{\frac{1}{n} \sum_{k=1}^n \sum_{\ell=1}^n \psi_{i_\ell}(x_k) - \sum_{\ell=1}^n \int \psi_{i_\ell} dx}$$

Clearly  $E_1 \tilde{f} = 0$ ,  $E_{v_n} \tilde{f} = 1$ . We have further

$$\text{VAR}_1 \tilde{f} = \|\tilde{f}\|_2^2 \leq \left( \frac{n\sqrt{\frac{1}{L} + \epsilon}}{1-n\epsilon - n\sqrt{\frac{1}{L} + \epsilon}} \right)^2$$

$$\text{VAR}_{v_n} \tilde{f} \leq \left( \frac{2n\epsilon}{1-n\epsilon - n\sqrt{\frac{1}{L} + \epsilon}} \right)^2$$

Hence,  $J_N^n(\tilde{f})$  becomes arbitrarily small as  $N \rightarrow \infty$  with probability arbitrarily close to one.

The solutions of (\*) can be easily obtained by the method of Lagrange multipliers. Since  $\varphi_1 = 1$ , (\*) is reduced to solving the following for  $a_i$  -

$$\min \left[ \frac{1}{2} \sum_{i=2}^N a_i^2 + \frac{1}{2} \sum_{i,j=2}^N a_i a_j \bar{\varphi}_{ij} \right]$$

such that

$$\sum_2^N a_i \bar{\varphi}_i = 1$$

$$\bar{\varphi}_i = \frac{1}{n} \sum_1^n \varphi_i(x_\ell)$$

$$\bar{\varphi}_{ij} = \frac{1}{n} \sum_1^n \varphi_i(x_\ell) \varphi_j(x_\ell)$$

For the determination of  $N=N_n$ , we first estimate  $J_{\frac{1}{2}}(\bar{f})$  by

$$\bar{J}_n = \frac{1}{2} + \frac{\epsilon^n}{4(1-\epsilon^n)} \quad (5)$$

where  $\epsilon^n$  is the expected nearest neighbor error rate of the  $n$  samples with the leaving-one-out method:

$$\epsilon^n = \frac{1}{n} \sum_{\ell=1}^n [1 - (1 - V_\ell)^{n-1}] \quad (6)$$

where  $V_\ell$  is the volume of the intersection of  $I$  with a sphere centered at  $x_\ell$  and of radius equal to the distance between  $x_\ell$  and its nearest neighbor in  $\{x_k\}_{k \neq \ell}$ . Now  $\epsilon^n \rightarrow \epsilon_{nn}$  with probability one as  $n \rightarrow \infty$  and hence  $\bar{J}_n \rightarrow J_{\frac{1}{2}}(\bar{f})$  with probability one as  $n \rightarrow \infty$ . Let  $N_n$  be an  $N$  which minimizes  $|J_N^n(f_n) - \bar{J}_n|$ . By the lemma such an  $N$  exists with probability one provided that  $\bar{J}_n > 0$  and this is true with probability one as  $n \rightarrow \infty$ .

The estimate we then use for  $p_2$  is

$$p_n = \frac{f_n - 1 + 2\bar{J}_n}{2\bar{J}_n - f_n} \quad (7)$$

If we should know the value of  $K$ , we may use the truncated estimate

$$\hat{p}_n = (p_n \vee 0) \wedge K \quad (8)$$

We now make the following consistency claim.

Theorem  $p_n \rightarrow p_2$  in Lebesgue measure with probability one and

$\hat{p}_n \xrightarrow{L_2} p_2$  with probability one.

Proof From the form of (3), (7), (8) and the fact that  $\bar{J}_n \rightarrow J_{\frac{1}{2}}(\bar{f})$ , it suffices to show that  $f_n \xrightarrow{L_2} \bar{f}$  with probability one.

Note that  $\varphi_2, \varphi_3, \dots$  are linearly independent and dense in  $\{f: \int f dx = 0\} \cap L_2(\frac{1}{2} + \frac{p_2}{2})$  where  $L_2(\frac{1}{2} + \frac{p_2}{2})$  denotes the set of square integrable functions wrt. to a measure whose density is  $\frac{1}{2} + p_2/2$ .

Now form a complete orthonormal basis  $\xi_2, \xi_3, \dots$  of

$\{f: \int f dx = 0\} \cap L_2(\frac{1}{2} + \frac{p_2}{2})$  where each  $\xi_i$  is a linear combination of  $\varphi_2, \varphi_3, \dots, \varphi_i$ . Let  $c_i = \int \xi_i p_2$ . Then  $\bar{f} = \sum_2^{\infty} b_i \xi_i$  where  $b_i$  is

the solution of  $\min \sum_2^{\infty} b_i^2$  such that  $\sum_2^{\infty} c_i b_i = 1$ . This is just  $b_i = c_i / \sum_2^{\infty} c_i^2$ .

Similarly, we form a complete orthonormal basis  $\eta_2^n, \eta_3^n, \dots$  of  $\{f: \int f=0\} \cap L_2(\frac{1}{2} + \frac{v_n}{2})$ , with each  $\eta_i^n$  a linear combination of  $\varphi_2, \varphi_3, \dots, \varphi_i$ . Let  $d_i^n = \int \eta_i^n v_n$ . The solution of (\*) is given by  $f_n = \sum_2^{N_n} d_i^n \eta_i^n / \sum_2^{N_n} (d_i^n)^2$ .

Clearly,  $d_i^n \rightarrow c_i$  and  $\|\eta_i^n - \xi_i\|_2 \rightarrow 0$  with probability one as  $n \rightarrow \infty$  for each  $i$ . Since

$$\left| \left( \sum_2^N c_i^2 \right)^{-1} - \left( \sum_2^N (d_i^n)^2 \right)^{-1} \right| \rightarrow 0$$

with probability one as  $n \rightarrow \infty$ , for each  $N$ , it follows that

$$\left| \left( \sum_2^{N_n} (d_i^n)^2 \right)^{-1} - \left( \sum_2^\infty c_i^2 \right)^{-1} \right| \rightarrow 0$$

with probability one as  $n \rightarrow \infty$ , and hence  $\sum_2^{N_n} (d_i^n)^2 \rightarrow \sum_2^\infty c_i^2$

with probability one as  $n \rightarrow \infty$ . Finally, for any  $\epsilon > 0$  pick  $M$

such that  $\sum_M^\infty c_i^2 < \epsilon$ . Then

$$\overline{\lim} \|f_n - \bar{f}\|_2 \leq \overline{\lim} \left\| \sum_M^{N_n} d_i^n \eta_i^n \right\|_2 + \left\| \sum_M^\infty c_i \xi_i \right\|_2$$

with probability one. But

$$\|g\|_2 \leq \sqrt{2} \|g\|_{\frac{1}{2} + \frac{p_2}{2}} \quad \text{and} \quad \|g\|_2 \leq \sqrt{2} \|g\|_{\frac{1}{2} + \frac{v_n}{2}}.$$

Hence  $\lim \|f_n - \bar{f}\|_2 \leq 2\sqrt{2\varepsilon}$  with probability one.

Since  $\varepsilon$  was arbitrary, the proof is complete.

#### IV. A REMARK ON HYPOTHESIS TESTING

If we are given two sets of data  $\{x_i\}_{i=1}^n$  and  $\{y_j\}_{j=1}^n$  and are then given the task of constructing an optimal discriminant between the two classes, we might solve

$$\min \frac{1}{2} \left[ \text{VA}_{\mu_n} + \frac{1}{2} \text{VAR}_{\nu_n} \right] \quad (**)$$

such that

$$E_{\mu_n} f = 0$$

$$E_{\nu_n} f = 1$$

$$f = \sum_{i=1}^{N_n} \varphi_i$$

$\mu_n, \nu_n$  empirical densities for  $\{x_i\}, \{y_j\}$

where  $N_n$  is determined by the analagous "stopping rule".

Unfortunately the consistency proof does not apply in this case since we are unable to find a bound for  $\|g\|_{\frac{p_1+p_2}{2}}$  in terms of  $\|g\|_{\frac{\nu_n+\mu_n}{2}}$ . Hence, the reference density  $p_1 \equiv 1$  "forced"

the consistency. We therefore recommend the estimates:  $\hat{p}_1$  by the method of III using  $\{x_i\}$  and then similarly  $\hat{p}_2$  using  $\{y_j\}$ . The discriminant  $\hat{p}_2/\hat{p}_1$  will then be optimal with probability one as  $n \rightarrow \infty$ .

#### REFERENCES

- [1]. L. Jones, " $K^{\text{th}}$  Order Solutions to the Problem of Finding Optimal Discriminant Functions", Technical Note 1980-4, Lincoln Laboratory, M.I.T. (10 January 1980), submitted to S.I.A.M. J. Appl. Math. DTIC AD-A0822396/3.
- [2]. T. Cover and P. Hart, "Nearest Neighbor Pattern Classification", IEEE Trans. Inf. Theory IT-13, pgs. 21-27 (1967).

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

(14) TN-1980-54

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER (17) ESD/IR-80-230	2. GOVT ACCESSION NO. AD A094728	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) (6) On Nonparametric Probability Density Estimation Using Orthogonal Series.		5. TYPE OF REPORT & PERIOD COVERED (9) Technical Note
7. AUTHOR(s) (10) Lee K. Jones		6. PERFORMING ORG. REPORT NUMBER Technical Note 1980-54
9. PERFORMING ORGANIZATION NAME AND ADDRESS Lincoln Laboratory, M.I.T. P.O. Box 73 Lexington, MA 02173 (16) 627A		8. CONTRACT OR GRANT NUMBER(s) (15) F19628-80-C-0002
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Systems Command, USAF Andrews AFB Washington, DC 20331		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Program Element No. 63311F Project No. 627A
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Electronic Systems Division Hanscom AFB Bedford, MA 01731 (12) 15		12. REPORT DATE (11) 26 November 1980
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		13. NUMBER OF PAGES 16
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15. SECURITY CLASS. (of this report) Unclassified
18. SUPPLEMENTARY NOTES  None		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  density estimation      Fourier coefficients      orthogonal expansions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  A method of density estimation is proposed, which is a rational modification of orthogonal expansions, combined with a stopping rule determined by a nearest neighbor statistic. This method yields consistent estimates and applies (in principle) to density estimation in any number of dimensions.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20/655 glu



